You have been scheduled for a flexible sigmoidoscopy

On	at	AM/PM at
	Folsom Sierra Endoscopy Center	
	1671 Creekside Drive, Suite 100	
	Folsom, CA 95630 (916) 983-7475	
	Please arrive 60 minutes early for registration	

IF YOU NEED TO RESCHEDULE OR CANCEL THIS PROCEDURE PLEASE CALL OUR OFFICE AT (916) 983-4444 OR (530) 622-6430

You will need to purchase 2 Fleet or Saline enemas at any pharmacy. Apply 1 enema 3 ho prior to the procedure and 1 enema one hour prior to the procedure.

You may be given medications before the procedure that will make you drowsy afterwards.

Someone you know must drive you home. Cab rides and public transportation are not acceptable.

You will be at the endoscopy center approximately two (2) hours.

Medications y	ou are taking
Aspirin:	Do not take any aspirin product fordays prior to your exam.
	Do not take any aspirin product on that day of exam unless instructed otherwise
Other anti-inf	lammatory drugs (i.e., Motrin, Advil, Aleve, Excedrine, Celebrex, Vioxx):
	Do not take these medications fordays prior to your exam.
	Do not take these medications on that day of exam unless instructed otherwise
Coumadin, W	arfarin or other blood thinning agents (i.e., Plavix, Ticlid, Lovenox, Aggrenox):
	Please notify your doctor if you are taking these types of medications
	STOP takingdays prior to the procedure
	You may take the last dose on
Diabetes med	licines:
	If you are on oral medications for diabetes, do not take the morning of the procedure
	You can resume this medication after the procedure when you are ready to eat.
	If you are taking insulin, please ask for detailed instructions.
You may take	all other usual daily medications on the day of the procedure.

No solid foods the entire day before the procedure <u>CLEAR LIQUIDS</u> only. You may have

clear liquids until 3 hours prior to the examination, then nothing by mouth.

Your follow-up visit with the Gastroenterologist is ______